

ISSN: 0974 - 0376

She Ecoscan : Special issue, Vol. IX: 727-730: 2016 AN INTERNATIONAL QUARTERLY JOURNAL OF ENVIRONMENTAL SCIENCES www.theecoscan.com

VITEX NEGUNDO: A GROWTH INHIBITORY AGENT AGAINST PROTEUS MIRABILIS

Ratna Sinha and M. P. Sinha

KEYWORDS

Proteus mirabilis Vitex negundo Agar disc diffusion method Alkaloids Tannins Saponins Flavonoids



Proceedings of National Conference on Harmony with Nature in Context of Resource Conservation and Climate Change (HARMONY - 2016) October 22 - 24, 2016, Hazaribag, organized by Department of Zoology, Botany, Biotechnology & Geology Vinoba Bhave University, Hazaribag (Jharkhand) 825301 in association with NATIONAL ENVIRONMENTALISTS ASSOCIATION, INDIA www.neaindia.org



RATNA SINHA^{1*} AND M. P. SINHA² ¹Department of Home Science, Ranchi University, Ranchi - 834 008 ²Department of Zoology, Ranchi University, Ranch - 834 008

e-mail: sinha.ratna20@gmail.com

ABSTRACT

In the present study the leaf extracts of Vitex negundo were analyzed for their phytochemical constituents and growth inhibitory impact against Proteus mirabilis by agar disc diffusion method. The phytochemcial screening revealed the presence of alkaloids, tannins, saponins and flavonoids. The concentration of tannins was highest and that of flavonoids was lowest. The inhibitory impact of the extract was measured as zone of inhibition (ZOI). The results were compared with standard antibiotic gentamycin. The study revealed that, the leaf extracts of Vitex negundo successfully inhibited the growth of Proteus mirabilis. The inhibitory impact of leaf extract of Vitex negundo against Proteus mirabilis seems to be due to presence of tannins, which are known for their antimicrobial impact. The antibacterial activity of Vitex negundo against P. mirabilis is indeed a promising development, which will help in the development of future antimicrobials.

INTRODUCTION

India is endowed with a rich wealth of medicinal plants. Herbs has always been playing chief role in medicinal formulations. The herbal medicines are gaining grounds throughout developing countries as people strive to stay healthy. India has recognized more than 2500 plant species which have medicinal values. However, large flora is still waiting for investigation for validation of their medicinal properties (Kirtikar and Basu, 1995). Vitex negundo belongs to family verbenaceae, known as Five leaf Chaste tree, is a large, aromatic shrub; with typical five foliate leaf pattern. It is found throughout the greater part of India at warmer zones and ascending to an altitude of 1500 m in outer western Himalayas. The shrub is one of the common plants used in Indian Medicines. It has been claimed to possess many medicinal properties. The plant has been reported to have high medicinal value (Rajith, 2011). Proteus mirabilis is a gram negative, facultatively anaerobic, rod shaped bacterium. It shows swarming motility and urease activity. Proteus mirabilis causes 90% of all infections in humans. It is widely distributed in soil and water. This rod shaped bacterium has the ability to produce high levels of urease, which hydrolyzes urea to ammonia (NH₂). Which makes the urine more alkaline. If left untreated the increased alkalinity can lead to formation of crystals or struvite, calcium carbonate, which can result in kidney stones. Proteus mirabilis is generally susceptible to most antibiotics apart from tetracycline and nitrofurantoin; 10 - 20 % of Proteus mirabilis strains are also resistant to first generation cephalosporins and ampicillin (O' Hara et al., 2000). The paper deals with growth inhibition of Proteus mirabilis by the methanolic extract of Vitex negundo.

MATERIALS AND METHODS

Plant Materials

The fresh tender leaves of *Vitex negundo* were collected and were washed with deionised water and disinfected with 0.1% HgCl₂ solution for 5 min and dried in shade for 15 days. The dried materials were ground to fine powder with help of electric grinder (Kumar et *al.*, 2013a; 2013b; 2013c).

Extract Preparation

50g of sieved powder was weighed and subjected to extraction in Soxhlet apparatus using methanol. The obtained extract was concentrated with the help of vacuum rotary evaporator and stored in air tight container (Kumar et al., 2013d; 2013e).

Phytochemical Screening

Qualitative and quantitative phytochemical screening was conducted as per previously published standards (Kumar et al., 2015a; 2015b).

Antibacterial Screening

The extract of Vitex negundo was used to screen the antibacterial action against Proteus mirabilis MTCC 1429

Agar Disc Diffusion Method

The agar disc diffusion method was used to analyse the antibacterial activity of

*Corresponding author



Figure 1: Showing the results of phytochemical screening of leaf extract of *Vitex negundo*



Figure 3: Showing Zone of Inhibition of gentamycin against *P. mirabilis*

Vitex negundo leaf extract agains *Proteus mirabilis*. Initially the stock cultures of bacteria were revived by inoculating in broth media and grown at 37° C for 18 hrs. the agar plate of above media were prepared and wells were made in the plate. Each plate was inoculated with 18h old cultures (100 μ L, 104 cfu) and spread evenly on the plate. After 20 min, the wells were filled with different concentrations of samples. The control wells were filled with gentamycin along with solvent. All the plates were incubated at 37°C for 24h and the diameter of inhibition zones were noted.

RESULTS AND DISCUSSION

Phytochemical Screening

The results of phytochemical screening of leaf extracts of *Vitex negundo* is presented as figure 1. The results reveal the presence of alkaloids, tannins, saponin, flavonoids. Tannins were found in highest concentration in leaf extract of *Vitex negundo*. Similar results were reported by Kumar et *al.*, (2013d). Several phytochemicals have been known to possess antibacterial properties. Tannins, alkaloids, saponins, flavonoids have been found to have growth inhibitory impact against *P. mirabilis*. Tannins have been reported to form irreversible complexes with prolene rich protein resulting in inhibition of cell synthesis of bacteria (Mamtha et *al.*, 2014)

Antibacterial analysis

The antibacterial analysis revealed that the leaf extract of Vitex



Figure 2: showing Zone of inhibition of Vitex negundo against P. mirabilis

negundo successfully inhibited the growth of *P. mirabilis*. The antibacterial activity is shown as agar plates in figure 2. The impact of standard antibiotic gentamycin against *P.* mirabilis is shown in figure 2. The extract showed a Zone of inhibition of 3, 4, 9 and 12 mm respectively for 0.612 mg, 1025 mg, 2.5 mg and 5 mg of extract concentration. Several phytochemicals have been known to possess antibacterial properties. Tannins, alkaloids, saponins, flavonoids have been found to growth inhibitory impact against *P. mirabilis* (Kumar et al., 2013d).

The antibacterial activity of *Vitex negundo* against *P. mirabilis* is indeed a promising development, which will help in development of future antimicrobial formulations against *Proteus mirabilis*.

REFERENCES

Kirtikar, K. R. and Basu, B. D. 1995. Indian medicinal plants. In International book distributors, Dehradun, 1: 830-832.

Kumar, M., Kumar, M., Dandapat, S. and Sinha, M. P. 2013a. Phytochemical screening and antioxidant potency of *Adhatoda vasica* and *Vitex negundo*. In *The Bioscan*. **8(2)**: 727-730.

Kumar, M., Dandapat, S., Kumar, A. and Sinha, M. P. 2013b. Determination of nutritive value and mineral elements of Five-leaf chaste tree (*Vitex negundo* L.) and Malabar Nut (*Adhatoda vasica* Nees). In *Academic J. Plant Sciences.* **6(3):** 103-108.

Kumar, M., Dandapat, S., Kumar, A. and Sinha, M. P.2013c. Pharmacological screening of leaf extract of *Adhatoda vasica* for therapeutic efficacy. In *Global J. Pharmacology*. **8(4)**: 494-500.

Kumar, M., Dandapat, S., Kumar, A. and Sinha, M. P. 2013d. Growth inhibitory impact of *Adhatoda vasica* and *Vitex negundo* on some human pathogens. In *The Ecoscan, Special issue*. (IV): 241-246.

Kumar, M., Dandapat, S., Kumar, A. and Sinha, M. P. 2013e. Antityphoid activity of Adhatoda vasica and Vitex negundo. In Persian *Culf Crop Protection*. 2(3): 64-75.

Kumar, M., Dandapat, S. and Sinha, M. P. 2015a. Phytochemical analyais and growth inhibitory impact of *Swertia chirayita* aqueous leaf extract against some human pathogens. In *World J. Zoology*. **10**(3): 188-190.

Kumar, M., Dandapat, S. and Sinha, M. P. 2015b. Hepatoprotective activity of *Adhatoda vasica* and *Vitex negundo* leaf extracts against carbon tetrachloride induced Hepatotoxicity in rats. In *Advances in Biological Research*. 9(4): 242-246.

Mamtha, B., Kavitha, K., Srinivasan, K. K. and Shivananda, P. G. 2004. An in-vitro study of the effect of *Centellaasiatica* (Indian Pennywort) on enteric pathogens. In *Indian J. Pharmacology*.

RATNA SINHA AND M. P. SINHA

36(1): 41.

O'Hara, C. M., Brenner, F. W. and Miller, J. M. 2000. Classification, identification and clinical significance of *Proteus*, *Providencia* and

Morganella. In Clin. Microbiol. Rev. 13(4): 534-546.

Rajith, R. 2011. Ethnomedicines of Kurichyas. Kanpur district, Kerala, India. In Indian J. Natural Product and Resources. 1(2): 149-153.